



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

XNI. *Experiments on Air.* By Henry Cavendish, Esq.
F. R. S. & S. A.

Read Jan. 15, 1784.

THE following experiments were made principally with a view to find out the cause of the diminution which common air is well known to suffer by all the various ways in which it is phlogistified, and to discover what becomes of the air thus lost or condensed; and as they seem not only to determine this point, but also to throw great light on the constitution and manner of production of dephlogistified air, I hope they may be not unworthy the acceptance of this society.

Many gentlemen have supposed that fixed air is either generated or separated from atmospheric air by phlogistification, and that the observed diminution is owing to this cause; my first experiments therefore were made in order to ascertain whether any fixed air is really produced thereby. Now, it must be observed, that as all animal and vegetable substances contain fixed air, and yield it by burning, distillation, or putrefaction, nothing can be concluded from experiments in which the air is phlogistified by them. The only methods I know, which are not liable to objection, are by the calcination of metals, the burning of sulphur or phosphorus, the mixture of nitrous air, and the explosion of inflammable air. Perhaps it may be supposed, that I ought to add to these the electric spark; but I think

think it much most likely, that the phlogification of the air, and production of fixed air, in this process, is owing to the burning of some inflammable matter in the apparatus. When the spark is taken from a solution of tournsol, the burning of the tournsol may produce this effect; when it is taken from lime-water, the burning of some foulness adhering to the tube, or perhaps of some inflammable matter contained in the lime, may have the same effect; and when quicksilver or metallic knobs are used, the calcination of them may contribute to the phlogification of the air, though not to the production of fixed air.

There is no reason to think that any fixed air is produced by the first method of phlogification. Dr. PRIESTLEY never found lime-water to become turbid by the calcination of metals over it*: Mr. LAVOISIER also found only a very slight and scarce perceptible turbid appearance, without any precipitation, to take place when lime-water was shaken in a glass vessel full of the air in which lead had been calcined; and even this small diminution of transparency in the lime-water might very likely arise, not from fixed air, but only from its being fouled by particles of the calcined metal, which we are told adhered in some places to the glass. This want of turbidity has been attributed to the fixed air uniting to the metallic calx, in preference to the lime; but there is no reason for supposing that the calx contained any fixed air; for I do not know that any one has extracted it from calces prepared in this manner; and though most metallic calces prepared over the fire, or by long exposure to the atmosphere, where they are in contact with fixed air, contain that substance, it by no means follows that they must

* Experiments on Air, vol. I. p. 137.

do so when prepared by methods in which they are not in contact with it.

Dr. PRIESTLEY also observed, that quicksilver, fouled by the addition of lead or tin, deposits a powder by agitation and exposure to the air, which consists in great measure of the calx of the imperfect metal. He found too some powder of this kind to contain fixed air *; but it is by no means clear that this air was produced by the phlogification of the air in which the quicksilver was shaken; as the powder was not prepared on purpose, but was procured from quicksilver fouled by having been used in various experiments, and may therefore have contained other impurities besides the metallic calces.

I never heard of any fixed air being produced by the burning of sulphur or phosphorus; but it has been asserted, and commonly believed, that lime water is rendered cloudy by a mixture of common and nitrous air; which, if true, would be a convincing proof that on mixing those two substances some fixed air is either generated or separated; I therefore examined this carefully. Now it must be observed, that as common air usually contains a little fixed air, which is no essential part of it, but is easily separated by lime water; and as nitrous air may also contain fixed air, either if the metal from which it is procured be rusty, or if the water of the vessel in which it is caught contain calcareous earth, suspended by fixed air, as most waters do, it is proper first to free both airs from it by previously washing them with lime water †. Now I found, by repeated

* *Exper. in Nat. Phil.* vol. I. p. 144.

† Though fixed air is absorbed in considerable quantity by water, as I shewed in *Phil. Transf.* vol. LVI. yet it is not easy to deprive common air of all the fixed

ed experiments, that if the lime water was clean, and the two airs were previously washed with that substance, not the least cloud was produced, either immediately on mixing them, or on suffering them to stand upwards of an hour, though it appeared by the thick clouds which were produced in the lime water, by breathing through it after the experiment was finished, that it was more than sufficient to saturate the acid formed by the decomposition of the nitrous air, and consequently that if any fixed air had been produced, it must have become visible. Once indeed I found a small cloud to be formed on the surface, after the mixture had stood a few minutes. In this experiment the lime water was not quite clean; but whether the cloud was owing to this circumstance, or to the air's having not been properly washed, I cannot pretend to say.

Neither does any fixed air seem to be produced by the explosion of the inflammable air obtained from metals, with either common or dephlogisticated air. This I tried by putting a little lime-water into a glass globe fitted with a brass cock, so as to make it air tight, and an apparatus for firing air by electricity. This globe was exhausted by an air-pump, and the two airs, which had been previously washed with lime-water, let in, and suffered to remain some time, to shew whether they would affect the lime-water, and then fired by electricity. The event was, that not the least cloud was produced in the lime-water, when the inflammable air was mixed with common air, and

air contained in it by means of water. On shaking a mixture of ten parts of common air, and one of fixed air, with more than an equal bulk of distilled water, not more than half of the fixed air was absorbed, and on transferring the air into fresh distilled water only half the remainder was absorbed, as appeared by the diminution which it still suffered on adding lime water.

only

only a very slight one, or rather diminution of transparency, when it was combined with dephlogisticated air. This, however, seemed not to be produced by fixed air; as it appeared instantly after the explosion, and did not increase on standing, and was spread uniformly through the liquor; whereas if it had been owing to fixed air, it would have taken up some short time before it appeared, and would have begun first at the surface, as was the case in the abovementioned experiment with nitrous air. What it was really owing to I cannot pretend to say; but if it did proceed from fixed air it would shew that only an excessively minute quantity was produced *. On the whole, though it is not improbable that fixed air may be generated in some chymical processes, yet it seems certain that it is not the general effect of phlogisticating air, and that the diminution of common air is by no means owing to the generation or separation of fixed air from it.

As there seemed great reason to think, from Dr. Priestley's experiments, that the nitrous and vitriolic acids were convertible into dephlogisticated air, I tried whether the dephlogisticated part of common air might not, by phlogistication, be changed into nitrous or vitriolic acid. For this purpose I impregnated some milk of lime with the fumes of burning sulphur, by putting a little of it into a large glass receiver, and burning sulphur therein, taking care to keep the mouth of the receiver stoppt till the fumes were all absorbed; after which the air of the receiver was changed, and more sulphur burnt in it as before, and the process repeated till 122 grains of sulphur were consumed. The milk of lime was then filtered and evaporated, but it yielded no nitrous salt, nor any other substance except selenite; so that no sensible quantity of the air was changed

* Dr. PRIESTLEY also found no fixed air to be produced by the explosion of inflammable and common air. Vol. V. p. 124.

into nitrous acid. It must be observed, that as the vitriolic acid produced by the burning sulphur is changed by its union with the lime into selenite, which is very little soluble in water, a very small quantity of nitrous salt, or any other substance which is soluble in water, would have been perceived.

I also tried whether any nitrous acid was produced by phlogisticating common air with liver of sulphur; for this purpose I made a solution of flowers of sulphur by boiling it with lime, and put a little of it into a large receiver, and shook it frequently, changing now and then the air, till the yellow colour of the solution was quite gone; a sign that all the sulphur was, by the loss of its phlogiston, turned into vitriolic acid, and united to the lime, or precipitated; the liquor was then filtered and evaporated, but it yielded not the least nitrous salt.

The experiment was repeated in nearly the same manner with dephlogisticated air procured from red precipitate; but not the least nitrous acid was obtained.

It is well known that common selenite is very little soluble in water; whereas that procured in the two last experiments was very soluble, and even crystallized readily, and was intensely bitter; this however appeared to be owing merely to the acid with which it was formed being very much phlogisticated; for on evaporating it to dryness, and exposing it to the air for a few days, it became much less soluble, so that on adding water to it not much dissolved; and by repeating this process once or twice, it seemed to become not more soluble than selenite made in the common manner.

This solubility of the selenite caused some trouble in trying the experiment; for while it continued much soluble it would have been impossible to have distinguished a small mixture of nitrous salt; but by the abovementioned process I was able to distinguish

distinguish as small a proportion as if the selenite had been originally no more soluble than usual.

The nature of the neutral salts made with the phlogisticated vitriolic and nitrous acids has not been much examined by the chymists, though it seems well worth their attention; and it is likely that many besides the foregoing may differ remarkably from those made with the same acids in their common state. Nitre formed with the phlogisticated nitrous acid has been found to differ considerably from common nitre, as well as Sal Polychrest from vitriolated tartar.

In order to try whether any vitriolic acid was produced by the phlogistication of air, I impregnated fifty ounces of distilled water with the fumes produced on mixing fifty-two ounce measures of nitrous air with a quantity of common air sufficient to decompose it. This was done by filling a bottle with some of this water, and inverting it into a basin of the same, and then, by a syphon, letting in as much nitrous air as filled it half-full; after which common air was added slowly by the same syphon, till all the nitrous air was decomposed. When this was done, the distilled water was further impregnated in the same manner till the whole of the abovementioned quantity of nitrous air was employed. This impregnated water, which was very sensibly acid to the taste, was distilled in a glass retort. The first runnings were very acid, and smelt pungent, being nitrous acid much phlogisticated; what came next had no sensible taste or smell; but the last runnings were very acid, and consisted of nitrous acid not phlogisticated. Scarce any sediment was left behind. These different parcels of distilled liquor were then exactly saturated with salt of tartar, and evaporated; they yielded $87\frac{1}{2}$ grains of nitre, which, as far as I could perceive, was unmixed with vitriolated tartar or any

other substance, and consequently no sensible quantity of the common air with which the nitrous air was mixed was turned into vitriolic acid.

It appears, from this experiment, that nitrous air contains as much acid as $2\frac{1}{4}$ times its weight of saltpetre; for fifty-two ounce measures of nitrous air weigh 32 grains, and, as was before said, yield as much acid as is contained in $87\frac{1}{2}$ grains of saltpetre; so that the acid in nitrous air is in a remarkably concentrated state, and I believe more than $1\frac{1}{2}$ times as much so as the strongest spirit of nitre ever prepared.

Having now mentioned the unsuccessful attempts I made to find out what becomes of the air lost by phlogification, I proceed to some experiments, which serve really to explain the matter.

In Dr. PRIESTLEY's last volume of experiments is related an experiment of Mr. WARLTIRE's, in which it is said that, on firing a mixture of common and inflammable air by electricity in a close copper vessel holding about three pints, a loss of weight was always perceived, on an average about two grains, though the vessel was stopped in such a manner that no air could escape by the explosion. It is also related, that on repeating the experiment in glass vessels, the inside of the glass, though clean and dry before, immediately became dewy; which confirmed an opinion he had long entertained, that common air deposits its moisture by phlogification. As the latter experiment seemed likely to throw great light on the subject I had in view, I thought it well worth examining more closely. The first experiment also, if there was no mistake in it, would be very extraordinary and curious; but it did not succeed with me; for though the vessel I used held more than Mr. WARLTIRE's, namely, 24,000 grains of water, and though the experiment

was

was repeated several times with different proportions of common and inflammable air, I could never perceive a loss of weight of more than one-fifth of a grain, and commonly none at all. It must be observed, however, that though there were some of the experiments in which it seemed to diminish a little in weight, there were none in which it increased*.

In all the experiments, the inside of the glass globe became dewy, as observed by Mr. WARLTIRE; but not the least footy matter could be perceived. Care was taken in all of them to find how much the air was diminished by the explosion, and to observe its test. The result is as follows: the bulk of the inflammable air being expressed in decimals of the common air,

Common air.	Inflammable air.	Diminution.	Air remaining after the explosion.	Test of this air in first method.	Standard.
1	1,241	,686	1,555	,055	,0
	1,055	,642	1,413	,063	,0
	,706	,647	1,059	,066	,0
	,423	,612	,811	,097	,03
	,331	,476	,855	,339	,27
	,206	,294	,912	,048	,58

In these experiments the inflammable air was procured from zinc, as it was in all my experiments, except where otherwise expressed: but I made two more experiments, to try whether there was any difference between the air from zinc and that from iron, the quantity of inflammable air being the same in both, namely. 0,331 of the common; but I could not find any difference to be depended on between the two kinds of air,

* Dr. PRIESTLEY, I am informed, has since found the experiment not to succeed.

either in the diminution which they suffered by the explosion, or the test of the burnt air.

From the fourth experiment it appears, that 423 measures of inflammable air are nearly sufficient to completely phlogificate 1000 of common air; and that the bulk of the air remaining after the explosion is then very little more than four-fifths of the common air employed; so that as common air cannot be reduced to a much less bulk than that by any method of phlogification, we may safely conclude, that when they are mixed in this proportion, and exploded, almost all the inflammable air, and about one-fifth part of the common air, lose their elasticity, and are condensed into the dew which lines the glasses.

The better to examine the nature of this dew, 500000 grain measures of inflammable air were burnt with about $2\frac{1}{2}$ times that quantity of common air, and the burnt air made to pass through a glass cylinder eight feet long and three-quarters of an inch in diameter, in order to deposit the dew. The two airs were conveyed slowly into this cylinder by separate copper pipes, passing through a brass plate which stopped up the end of the cylinder; and as neither inflammable nor common air can burn by themselves, there was no danger of the flame spreading into the magazines from which they were conveyed. Each of these magazines consisted of a large tin vessel, inverted into another vessel just big enough to receive it. The inner vessel communicated with the copper pipe, and the air was forced out of it by pouring water into the outer vessel; and in order that the quantity of common air expelled should be $2\frac{1}{2}$ times that of the inflammable, the water was let into the outer vessels by two holes in the bottom of the same tin pan, the hole which conveyed the water into that vessel in
which

which the common air was confined being $2\frac{1}{2}$ times as big as the other.

In trying the experiment, the magazines being first filled with their respective airs, the glass cylinder was taken off, and water let, by the two holes, into the outer vessels, till the airs began to issue from the ends of the copper pipes; they were then set on fire by a candle, and the cylinder put on again in its place. By this means upwards of 135 grains of water were condensed in the cylinder, which had no taste nor smell, and which left no sensible sediment when evaporated to dryness; neither did it yield any pungent smell during the evaporation; in short, it seemed pure water.

In my first experiment, the cylinder near that part where the air was fired was a little tinged with sooty matter, but very slightly so; and that little seemed to proceed from the putty with which the apparatus was luted, and which was heated by the flame; for in another experiment, in which it was contrived so that the luting should not be much heated, scarce any sooty tinge could be perceived.

By the experiments with the globe it appeared, that when inflammable and common air are exploded in a proper proportion, almost all the inflammable air, and near one-fifth of the common air, lose their elasticity, and are condensed into dew. And by this experiment it appears, that this dew is plain water, and consequently that almost all the inflammable air, and about one-fifth of the common air, are turned into pure water.

In order to examine the nature of the matter condensed on firing a mixture of dephlogisticated and inflammable air, I took a glass globe, holding 8800 grain measures, furnished with a brass cock and an apparatus for firing air by electricity. This globe was well exhausted by an air-pump, and then filled with

a mixture of inflammable and dephlogisticated air, by shutting the cock, fastening a bent glass tube to its mouth, and letting up the end of it into a glass jar inverted into water, and containing a mixture of 19500 grain measures of dephlogisticated air, and 37000 of inflammable; so that, upon opening the cock, some of this mixed air rushed through the bent tube, and filled the globe*. The cock was then shut, and the included air fired by electricity, by which means almost all of it lost its elasticity. The cock was then again opened, so as to let in more of the same air, to supply the place of that destroyed by the explosion, which was again fired, and the operation continued till almost the whole of the mixture was let into the globe and exploded. By this means, though the globe held not more than the sixth part of the mixture, almost the whole of it was exploded therein, without any fresh exhaustion of the globe.

As I was desirous to try the quantity and test of this burnt air, without letting any water into the globe, which would have prevented my examining the nature of the condensed matter, I took a larger globe, furnished also with a stop cock, exhausted it by an air-pump, and screwed it on upon the cock of the former globe; upon which, by opening both cocks, the air rushed out of the smaller globe into the larger, till it became of equal density in both; then, by shutting the cock of the larger globe, unscrewing it again from the former, and opening it under water, I was enabled to find the quantity of the burnt air in it; and consequently, as the proportion which the contents of the two globes bore to each other was

* In order to prevent any water from getting into this tube, while dipped under water to let it up into the glass jar, a bit of wax was stuck upon the end of it, which was rubbed off when raised above the surface of the water.

known, could tell the quantity of burnt air in the small globe before the communication was made between them. By this means the whole quantity of the burnt air was found to be 2950 grain measures; its standard was 1,85.

The liquor condensed in the globe, in weight about 30 grains, was sensibly acid to the taste, and by saturation with fixed alkali, and evaporation, yielded near two grains of nitre; so that it consisted of water united to a small quantity of nitrous acid. No sooty matter was deposited in the globe. The dephlogisticated air used in this experiment was procured from red precipitate, that is, from a solution of quicksilver in spirit of nitre distilled till it acquires a red colour.

As it was suspected, that the acid contained in the condensed liquor was no essential part of the dephlogisticated air, but was owing to some acid vapour which came over in making it and had not been absorbed by the water, the experiment was repeated in the same manner, with some more of the same air, which had been previously washed with water, by keeping it a day or two in a bottle with some water, and shaking it frequently; whereas that used in the preceding experiment had never passed through water, except in preparing it. The condensed liquor was still acid.

The experiment was also repeated with dephlogisticated air, procured from red lead by means of oil of vitriol; the liquor condensed was acid, but by an accident I was prevented from determining the nature of the acid.

I also procured some dephlogisticated air from the leaves of plants, in the manner of Doctors INGENHOUSZ and PRIESTLEY, and exploded it with inflammable air as before; the condensed liquor still continued acid, and of the nitrous kind.

In all these experiments the proportion of inflammable air was such, that the burnt air was not much phlogificated; and it was observed, that the less phlogificated it was, the more acid was the condensed liquor. I therefore made another experiment, with some more of the same air from plants, in which the proportion of inflammable air was greater, so that the burnt air was almost completely phlogificated, its standard being $\frac{1}{10}$. The condensed liquor was then not at all acid, but seemed pure water: so that it appears, that with this kind of dephlogificated air, the condensed liquor is not at all acid, when the two airs are mixed in such a proportion that the burnt air is almost completely phlogificated, but is considerably so when it is not much phlogificated.

In order to see whether the same thing would obtain with air procured from red precipitate, I made two more experiments with that kind of air, the air in both being taken from the same bottle, and the experiment tried in the same manner, except that the proportions of inflammable air were different. In the first, in which the burnt air was almost completely phlogificated, the condensed liquor was not at all acid. In the second, in which its standard was 1,86, that is, not much phlogificated, it was considerably acid; so that with this air, as well as with that from plants, the condensed liquor contains, or is entirely free from, acid, according as the burnt air is less or more phlogificated; and there can be little doubt but that the same rule obtains with any other kind of dephlogificated air.

In order to see whether the acid, formed by the explosion of dephlogificated air obtained by means of the vitriolic acid, would also be of the nitrous kind, I procured some air from turbith mineral, and exploded it with inflammable air, the
 propor-

proportion being such that the burnt air was not much phlogisticated. The condensed liquor manifested an acidity, which appeared, by saturation with a solution of salt of tartar, to be of the nitrous kind; and it was found, by the addition of some terra ponderosa salita, to contain little or no vitriolic acid.

When inflammable air was exploded with common air, in such a proportion that the standard of the burnt air was about $\frac{4}{10}$, the condensed liquor was not in the least acid. There is no difference, however, in this respect between common air, and dephlogisticated air mixed with phlogisticated in such a proportion as to reduce it to the standard of common air; for some dephlogisticated air from red precipitate, being reduced to this standard by the addition of perfectly phlogisticated air, and then exploded with the same proportion of inflammable air as the common air was in the foregoing experiment, the condensed liquor was not in the least acid.

From the foregoing experiments it appears, that when a mixture of inflammable and dephlogisticated air is exploded in such proportion that the burnt air is not much phlogisticated, the condensed liquor contains a little acid, which is always of the nitrous kind, whatever substance the dephlogisticated air is procured from; but if the proportion be such that the burnt air is almost entirely phlogisticated, the condensed liquor is not at all acid, but seems pure water, without any addition whatever; and as, when they are mixed in that proportion, very little air remains after the explosion, almost the whole being condensed, it follows, that almost the whole of the inflammable and dephlogisticated air is converted into pure water. It is not easy, indeed, to determine from these experiments what proportion the burnt air, remaining after the explosions, bore to the dephlogisticated air employed, as neither the

small nor the large globe could be perfectly exhausted of air, and there was no saying with exactness what quantity was left in them; but in most of them, after allowing for this uncertainty, the true quantity of burnt air seemed not more than $\frac{1}{17}$ th of the dephlogisticated air employed, or $\frac{1}{10}$ th of the mixture. It seems, however, unnecessary to determine this point exactly, as the quantity is so small, that there can be little doubt but that it proceeds only from the impurities mixed with the dephlogisticated and inflammable air, and consequently that, if those airs could be obtained perfectly pure, the whole would be condensed.

With respect to common air, and dephlogisticated air reduced by the addition of phlogisticated air to the standard of common air, the case is different; as the liquor condensed in exploding them with inflammable air, I believe I may say in any proportion, is not at all acid; perhaps, because if they are mixed in such a proportion as that the burnt air is not much phlogisticated, the explosion is too weak, and not accompanied with sufficient heat.

All the foregoing experiments, on the explosion of inflammable air with common and dephlogisticated airs, except those which relate to the cause of the acid found in the water, were made in the summer of the year 1781, and were mentioned by me to Dr. PRIESTLEY, who in consequence of it made some experiments of the same kind, as he relates in a paper printed in the preceding volume of the Transactions. During the last summer also, a friend of mine gave some account of them to M. LAVOISIER, as well as of the conclusion drawn from them, that dephlogisticated air is only water deprived of phlogiston; but at that time so far was M. LAVOISIER from thinking any such opinion warranted, that, till he was pre-
vailed

vailed upon to repeat the experiment himself, he found some difficulty in believing that nearly the whole of the two airs could be converted into water. It is remarkable, that neither of these gentlemen found any acid in the water produced by the combustion ; which might proceed from the latter having burnt the two airs in a different manner from what I did ; and from the former having used a different kind of inflammable air, namely, that from charcoal, and perhaps having used a greater proportion of it.

Before I enter into the cause of these phænomena, it will be proper to take notice, that phlogisticated air appears to be nothing else than the nitrous acid united to phlogiston ; for when nitre is deflagrated with charcoal, the acid is almost entirely converted into this kind of air. That the acid is entirely converted into air, appears from the common process for making what is called clyffus of nitre ; for if the nitre and charcoal are dry, scarce any thing is found in the vessels prepared for condensing the fumes ; but if they are moist a little liquor is collected, which is nothing but the water contained in the materials, impregnated with a little volatile alkali, proceeding in all probability from the imperfectly burnt charcoal, and a little fixed alkali, consisting of some of the alkalized nitre carried over by the heat and watery vapours. As far as I can perceive too, at present, the air into which much the greatest part of the acid is converted, differs in no respect from common air phlogisticated. A small part of the acid, however, is turned into nitrous air, and the whole is mixed with a good deal of fixed, and perhaps a little inflammable air, both proceeding from the charcoal.

It is well known, that the nitrous acid is also converted by phlogistication into nitrous air, in which respect there seems a considerable

considerable analogy between that and the vitriolic acid; for the vitriolic acid, when united to a smaller proportion of phlogiston, forms the volatile sulphureous acid and vitriolic acid air, both of which, by exposure to the atmosphere, lose their phlogiston, though not very fast, and are turned back into vitriolic acid; but, when united to a greater proportion of phlogiston, it forms sulphur, which shews no signs of acidity, unless a small degree of affinity to alkalies can be called so, and in which the phlogiston is more strongly adherent, so that it does not fly off when exposed to the air, unless assisted by a heat sufficient to set it on fire. In like manner the nitrous acid, united to a certain quantity of phlogiston, forms nitrous fumes and nitrous air, which readily quit their phlogiston to common air; but when united to a different, in all probability a larger quantity, it forms phlogisticated air, which shews no signs of acidity, and is still less disposed to part with its phlogiston than sulphur.

This being premised, there seem two ways by which the phenomena of the acid found in the condensed liquor may be explained; first, by supposing that dephlogisticated air contains a little nitrous acid which enters into it as one of its component parts, and that this acid, when the inflammable air is in a sufficient proportion, unites to the phlogiston, and is turned into phlogisticated air, but does not when the inflammable air is in too small a proportion; and, secondly, by supposing that there is no nitrous acid mixed with, or entering into the composition of, dephlogisticated air, but that, when this air is in a sufficient proportion, part of the phlogisticated air with which it is debased is, by the strong affinity of phlogiston to dephlogisticated air, deprived of its phlogiston and turned into nitrous acid; whereas, when the dephlogisticated air is not more than sufficient to consume the inflammable air,

none

none then remains to deprive the phlogificated air of its phlogiston, and turn it into acid.

If the latter explanation be true, I think, we must allow that dephlogificated air is in reality nothing but dephlogificated water, or water deprived of its phlogiston; or, in other words, that water consists of dephlogificated air united to phlogiston; and that inflammable air is either pure phlogiston, as Dr. PRIESTLEY and Mr. KIRWAN suppose, or else water united to phlogiston*; since, according to this supposition, these two substances united together form pure water. On the other hand, if the first explanation be true, we must suppose that dephlogificated air consists of water united to a little nitrous acid and deprived of its phlogiston; but still the nitrous acid in it must make only a very small part of the whole,

* Either of these suppositions will agree equally well with the following experiments; but the latter seems to me much the most likely. What principally makes me think so is, that common or dephlogificated air do not absorb phlogiston from inflammable air, unless assisted by a red heat, whereas they absorb the phlogiston of nitrous air, liver of sulphur, and many other substances, without that assistance; and it seems inexplicable, that they should refuse to unite to pure phlogiston, when they are able to extract it from substances to which it has an affinity; that is, that they should overcome the affinity of phlogiston to other substances, and extract it from them, when they will not even unite to it when presented to them. On the other hand, I know no experiment which shews inflammable air to be pure phlogiston rather than an union of it with water, unless it be Dr. PRIESTLEY's experiment of expelling inflammable air from iron by heat alone. I am not sufficiently acquainted with the circumstances of that experiment to argue with certainty about it; but I think it much more likely, that the inflammable air was formed by the union of the phlogiston of the iron filings with the water dispersed among them, or contained in the retort or other vessel in which it was heated; and in all probability this was the cause of the separation of the phlogiston, as iron seems not disposed to part with its phlogiston by heat alone, without being assisted by the air or some other substance.

as it is found, that the phlogisticated air, which it is converted into, is very small in comparison of the dephlogisticated air.

I think the second of these explanations seems much the most likely; as it was found, that the acid in the condensed liquor was of the nitrous kind, not only when the dephlogisticated air was prepared from red precipitate, but also when it was procured from plants or from turbith mineral: and it seems not likely, that air procured from plants, and still less likely that air procured from a solution of mercury in oil of vitriol, should contain any nitrous acid.

Another strong argument in favour of this opinion is, that dephlogisticated air yields no nitrous acid when phlogisticated by liver of sulphur; for if this air contains nitrous acid, and yields it when phlogisticated by explosion with inflammable air, it is very extraordinary that it should not do so when phlogisticated by other means.

But what forms a stronger and, I think, almost decisive argument in favour of this explanation is, that when the dephlogisticated air is very pure, the condensed liquor is made much more strongly acid by mixing the air to be exploded with a little phlogisticated air, as appears by the following experiments.

A mixture of 18500 grain measures of inflammable air with 9750 of dephlogisticated air procured from red precipitate were exploded in the usual manner; after which, a mixture of the same quantities of the same dephlogisticated and inflammable air, with the addition of 2500 of air phlogisticated by iron filings and sulphur, was treated in the same manner. The condensed liquor, in both experiments, was acid, but that in the latter evidently more so, as appeared also by saturating each of them separately with marble powder, and precipitating

the earth by fixed alkali, the precipitate of the second experiment weighing one-fifth of a grain, and that of the first being several times less. The standard of the burnt air in the first experiment was 1,86, and in the second only 0,9.

It must be observed, that all circumstances were the same in these two experiments, except that in the latter the air to be exploded was mixed with some phlogificated air, and that in consequence the burnt air was more phlogificated than in the former; and from what has been before said, it appears, that this latter circumstance ought rather to have made the condensed liquor less acid; and yet it was found to be much more so, which shews strongly that it was the phlogificated air which furnished the acid.

As a further confirmation of this point, these two comparative experiments were repeated with a little variation, namely, in the first experiment there was first let into the globe 1500 of dephlogificated air, and then the mixture, consisting of 12200 of dephlogificated air and 25900 of inflammable, was let in at different times as usual. In the second experiment, besides the 1500 of dephlogificated air first let in, there was also admitted 2500 of phlogificated air, after which the mixture, consisting of the same quantities of dephlogificated and inflammable air as before, was let in as usual. The condensed liquor of the second experiment was about three times as acid as that of the first, as it required 119 grains of a diluted solution of salt of tartar to saturate it, and the other only 37. The standard of the burnt air was 0,78 in the second experiment, and 1,96 in the first.

The intention of previously letting in some dephlogificated air in the two last experiments was, that the condensed liquor

was expected to become more acid thereby, as proved actually to be the case.

In the first of these two experiments, in order that the air to be exploded should be as free as possible from common air, the globe was first filled with a mixture of dephlogisticated and inflammable air, it was then exhausted, and the air to be exploded let in; by which means, though the globe was not perfectly exhausted, very little common air could be left in it. In the first set of experiments this circumstance was not attended to, and the purity of the dephlogisticated air was forgot to be examined in both sets.

From what has been said there seems the utmost reason to think, that dephlogisticated air is only water deprived of its phlogiston, and that inflammable air, as was before said, is either phlogisticated water, or else pure phlogiston; but in all probability the former.

As Mr. WATT, in a paper lately read before this Society, supposes water to consist of dephlogisticated air and phlogiston deprived of part of their latent heat, whereas I take no notice of the latter circumstance, it may be proper to mention in a few words the reason of this apparent difference between us. If there be any such thing as elementary heat, it must be allowed that what Mr. WATT says is true; but by the same rule we ought to say, that the diluted mineral acids consist of the concentrated acids united to water and deprived of part of their latent heat; that solutions of sal ammoniac, and most other other neutral salts, consist of the salt united to water and elementary heat; and a similar language ought to be used in speaking of almost all chemical combinations, as there are very few which are not attended with some increase or diminution of heat. Now I have chosen to avoid this form of speaking,
both

Both because I think it more likely that there is no such thing as elementary heat, and because saying so in this instance, without using similar expressions in speaking of other chemical unions, would be improper, and would lead to false ideas; and it may even admit of doubt, whether the doing it in general would not cause more trouble and perplexity than it is worth.

There is the utmost reason to think, that dephlogisticated and phlogisticated air, as M. LAVOISIER and SCHEELÉ suppose, are quite distinct substances, and not differing only in their degree of phlogistication; and that common air is a mixture of the two; for if the dephlogisticated air is pretty pure, almost the whole of it loses its elasticity by phlogistication, and, as appears by the foregoing experiments, is turned into water, instead of being converted into phlogisticated air. In most of the foregoing experiments, at least $\frac{1}{7}$ ths of the whole was turned into water; and by treating some dephlogisticated air with liver of sulphur, I have reduced it to less than $\frac{1}{30}$ th of its original bulk, and other persons, I believe, have reduced it to a still less bulk; so that there seems the utmost reason to suppose, that the small residuum which remains after its phlogistication proceeds only from the impurities mixed with it.

It was just said, that some dephlogisticated air was reduced by liver of sulphur to $\frac{1}{30}$ th of its original bulk; the standard of this air was 4,8, and consequently the standard of perfectly pure dephlogisticated air should be very nearly 5, which is a confirmation of the foregoing opinion; for if the standard of pure dephlogisticated air is 5, common air must, according to this opinion, contain one-fifth of it, and therefore ought to lose one-fifth of its bulk by phlogistication, which is what it is actually found to lose.

From

From what has been said, it follows, that instead of saying air is phlogisticated or dephlogisticated by any means, it would be more strictly just to say, it is deprived of, or receives, an addition of dephlogisticated air; but as the other expression is convenient, and can scarcely be considered as improper, I shall still frequently make use of it in the remainder of this paper.

There seemed great reason to think, from Dr. PRIESTLEY's experiments, that both the nitrous and vitriolic acids were convertible into dephlogisticated air, as that air is procured in the greatest quantity from substances containing those acids, especially the former. The foregoing experiments, however, seem to shew that no part of the acid is converted into dephlogisticated air, and that their use in preparing it is owing only to the great power which they possess of depriving bodies of their phlogiston. A strong confirmation of this is, that red precipitate, which is one of the substances yielding dephlogisticated air in the greatest quantity, and which is prepared by means of the nitrous acid, contains in reality no acid. This I found by grinding 400 grains of it with spirits of sal ammoniac, and keeping them together for some days in a bottle, taking care to shake them frequently. The red colour of the precipitate was rendered pale, but not entirely destroyed; being then washed with water and filtered, the clear liquor yielded on evaporation not the least ammoniacal salt.

It is natural to think, that if any nitrous acid had been contained in the red precipitate, it would have united to the volatile alkali and have formed ammoniacal nitre, and would have been perceived on evaporation; but in order to determine more certainly whether this would be the case, I dried some of the same solution of quicksilver from which the red precipitate was prepared with a less heat, so that it acquired only an orange colour,

colour, and treated the same quantity of it with volatile alkali in the same manner as before. It immediately caused an effervescence, changed the colour to grey, and yielded 52 grains of ammoniacal nitre. There is the utmost reason to think, therefore, that red precipitate contains no nitrous acid; and consequently that, in procuring dephlogisticated air from it, no acid is converted into air; and it is reasonable to conclude, therefore, that no such change is produced in procuring it from any other substance.

It remains to consider in what manner these acids act in producing dephlogisticated air. The way in which the nitrous acid acts, in the production of it from red precipitate, seems to be as follows. On distilling the mixture of quicksilver and spirit of nitre, the acid comes over, loaded with phlogiston, in the form of nitrous vapour, and continues to do so till the remaining matter acquires its full red colour, by which time all the nitrous acid is driven over, but some of the watery part still remains behind, and adheres strongly to the quicksilver; so that the red precipitate may be considered, either as quicksilver deprived of part of its phlogiston, and united to a certain portion of water, or as quicksilver united to dephlogisticated air*; after which, on further increasing the heat, the water in it rises deprived of its phlogiston, that is, in the form of dephlogisticated

* Unless we were much better acquainted than we are with the manner in which different substances are united together in compound bodies, it would be ridiculous to say, that it is the quicksilver in the red precipitate which is deprived of its phlogiston, and not the water, or that it is the water and not the quicksilver; all that we can say is, that red precipitate consists of quicksilver and water, one or both of which are deprived of part of their phlogiston. In like manner, during the preparation of the red precipitate, it is certain that the acid absorbs phlogiston, either from the quicksilver or the water; but we are by no means authorised to say from which.

air,

air, and at the same time the quicksilver distils over in its metallic form. It is justly remarked by Dr. PRIESTLEY, that the solution of quicksilver does not begin to yield dephlogisticated air till it acquires its red colour.

Mercurius calcinatus appears to be only quicksilver which has absorbed dephlogisticated air from the atmosphere during its preparation; accordingly, by giving it a sufficient heat, the dephlogisticated air is driven off, and the quicksilver acquires its original form. It seems therefore that mercurius calcinatus and red precipitate, though prepared in a different manner, are very nearly the same thing.

From what has been said it follows, that red precipitate and mercurius calcinatus contain as much phlogiston as the quicksilver they are prepared from; but yet, as uniting dephlogisticated air to a metal comes to the same thing as depriving it of part of its phlogiston and adding water to it, the quicksilver may still be considered as deprived of its phlogiston; but the imperfect metals seem not only to absorb dephlogisticated air during their calcination, but also to be really deprived of part of their phlogiston, as they do not acquire their metallic form by driving off the dephlogisticated air.

In procuring dephlogisticated air from nitre, the acid acts in a different manner, as, upon heating the nitre red-hot, the dephlogisticated air rises mixed with a little nitrous acid, and at the same time the acid remaining in the nitre becomes very much phlogisticated; which shews that the acid absorbs phlogiston from the water in the nitre, and becomes phlogisticated, while the water is thereby turned into dephlogisticated air. On distilling 3155 grains of nitre in an unglazed earthen retort, it yielded 256000 grain measures of dephlogisticated air*, the
standard

* This is, about eighty-one grain measures from one grain of nitre; and the weight

standard of different parts of which varied from 3 to 3,65, but at a medium was 3,35. The matter remaining in the retort dissolved readily in water, and tasted alkaline and caustic. On adding diluted spirit of nitre to the solution, strong red fumes were produced; a sign that the acid in it was very much phlogisticated, as no fumes whatever would have been produced on adding the same acid to a solution of common nitre; that part of the solution also which was supersaturated with acid became blue; a colour which the diluted nitrous acid is known to assume when much phlogisticated. The solution, when saturated with this acid, lost its alkaline and caustic taste, but yet tasted very different from true nitre, seeming as if it had been mixed with sea-salt, and also required much less water to dissolve it; but on exposing it for some days to the air, and adding fresh acid as fast as by the flying off of the fumes the alkali predominated, it became true nitre, unmixed, as far as I could perceive, with any other salt*.

It has been remarked, that the dephlogisticated air procured from nitre is less pure, than that from red precipitate and many other substances, which may perhaps proceed from unglazed earthen retorts having been commonly used for this purpose, and which, conformably to Dr. PRIESTLEY's discovery, may possibly absorb some common air from without, and emit it along with the dephlogisticated air; but if it should be found that the dephlogisticated air procured from nitre in glass or glazed earthen vessels is also impure, it would seem to shew that part

weight of the dephlogisticated air, supposing it 800 times lighter than water, is one tenth of that of the nitre. In all probability it would have yielded a much greater quantity of air, if a greater heat had been applied.

* This phlogistication of the acid in nitre by heat has been observed by Mr. SCHEELE; see his experiments on air and fire, p. 45. English translation.

of the acid in the nitre is turned into phlogisticated air, by absorbing phlogiston from the watery part.

From what has been said it appears, that there is a considerable difference in the manner in which the acid acts in the production of dephlogisticated air from red precipitate and from nitre; in the former case the acid comes over first, leaving the remaining substance deprived of part of its phlogiston; in the latter the dephlogisticated air comes first, leaving the acid loaded with the phlogiston of the water from which it was formed.

On distilling a mixture of quicksilver and oil of vitriol to dryness, part of the acid comes over, loaded with phlogiston, in the form of volatile sulphureous acid and vitriolic acid air; so that the remaining white mass may be considered as consisting of quicksilver deprived of its phlogiston, and united to a certain proportion of acid and water, or of plain quicksilver united to a certain proportion of acid and dephlogisticated air. Accordingly on urging this white mass with a more violent heat, the dephlogisticated air comes over, and at the same time part of the quicksilver rises in its metallic form, and also part of the white mass, united in all probability to a greater proportion of acid than before, sublimes; so that the rationale of the production of dephlogisticated air from turbith mineral, and from red precipitate, are nearly similar.

True turbith mineral consists of the abovementioned white mass, well washed with water, by which means it acquires a yellow colour, and contains much less acid than the unwashed mass. Accordingly it seems likely, that on exposing this to heat, less of it should sublime without being decomposed, and consequently that more dephlogisticated air should be procured from it than from the unwashed mass.

This

This is an instance, that the superabundant vitriolic acid may, in some cases, be better extracted from the base it is united to by water than by heat. Vitriolated tartar is another instance; for, if vitriolated tartar be mixed with oil of vitriol and exposed even to a pretty strong red heat, the mass will be very acid; but, if this mass is dissolved in water, and evaporated, the crystals will be not sensibly so.

In all probability, the vitriolic acid acts in the same manner in the production of dephlogisticated air from alum, as the nitrous does in its production from nitre; that is, the watery part comes over first in the form of dephlogisticated air, leaving the acid charged with its phlogiston. Whether this is also the case with regard to green and blue vitriol, or whether in them the acid does not rather act in the same manner as in turbith mineral, I cannot pretend to say, but I think the latter more likely.

There is another way by which dephlogisticated air has been found to be produced in great quantities, namely, the growth of vegetables exposed to the sun or day-light; the rationale of which, in all probability, is, that plants, when assisted by the light, deprive part of the water sucked up by their roots of its phlogiston, and turn it into dephlogisticated air, while the phlogiston unites to, and forms part of, the substance of the plant.

There are many circumstances which shew, that light has a remarkable power in enabling one body to absorb phlogiston from another. Mr. SENEBIER has observed, that the green tincture procured from the leaves of vegetables by spirit of wine, quickly loses its colour when exposed to the sun in a bottle not more than one third part full, but does not do so in the dark, or if the bottle is quite full of the tincture, or if the air in it

is phlogificated; whence it is natural to conclude, that the light enables the dephlogificated part of the air to absorb phlogiston from the tincture; and this appears to be really the case, as I find that the air in the bottle is considerably phlogificated thereby. Dephlogificated spirit of nitre also acquires a yellow colour, and becomes phlogificated, by exposure to the sun's rays *; and I find on trial that the air in the bottle in which it is contained becomes dephlogificated, or, in other words, receives an increase of dephlogificated air, which shews that the change in the acid is not owing to the sun's rays communicating phlogiston to it, but to their enabling it to absorb phlogiston from the water contained in it, and thereby to produce dephlogificated air. Mr. SCHEELÉ also found, that the dark colour acquired by luna cornea on exposure to the light, is owing to part of the silver being revived; and that gold, dissolved in aqua regia and deprived by distillation of the nitrous and superfluous marine acid, is revived by the same means; and there is the utmost reason to think, that, in both cases, the revival of the metal is owing to its absorbing phlogiston from the water.

Vegetables seem to consist almost intirely of fixed and phlogificated air, united to a large proportion of phlogiston and some water, since by burning in the open air, in which their phlogiston unites to the dephlogificated part of the atmosphere and forms

* If spirit of nitre is distilled with a very gentle heat, the part which comes over is high coloured and fuming, and that which remains behind is quite colourless, and fumes much less than other nitrous acid of the same strength, and the fumes are colourless. This is called dephlogificated spirit of nitre, as it appears to be really deprived of phlogiston by the process. The manner of preparing it, as well as its property of regaining its yellow colour by exposure to the light, is mentioned by Mr. SCHEELÉ in the Stockholm Memoirs, 1774.

water,

water, they seem to be reduced almost intirely to water and those two kinds of air. Now plants growing in water without earth, can receive nourishment only from the water and air, and must therefore in all probability absorb their phlogiston from the water. It is known also that plants growing in the dark do not thrive well, and grow in a very different manner from what they do when exposed to the light.

From what has been said, it seems likely that the use of light, in promoting the growth of plants and the production of dephlogisticated air from them, is, that it enables them to absorb phlogiston from the water. To this it may perhaps be objected, that though plants do not thrive well in the dark, yet they do grow, and should therefore, according to this hypothesis, absorb water from the atmosphere, and yield dephlogisticated air, which they have not been found to do. But we have no proof that they grew at all in any of those cases in which they were found not to yield dephlogisticated air; for though they will grow in the dark, yet their vegetative powers may perhaps at first be intirely checked by it, especially considering the unnatural situation in which they must be placed in such experiments. Perhaps too plants growing in the dark may be able to absorb phlogiston from water not much impregnated with dephlogisticated air, but not from water strongly impregnated with it; and consequently, when kept under water in the dark, may perhaps at first yield some dephlogisticated air, which, instead of rising to the surface, may be absorbed by the water, and, before the water is so much impregnated as to suffer any to escape, the plant may cease to vegetate, unless the water is changed. Unless therefore it could be shewn that plants growing in the dark, in water alone, will increase in size, without yielding dephlogisticated

gified air, and without the water becoming more impregnated with it than before, no objection can be drawn from thence.

Mr. SENEBIER finds, that plants yield much more dephlogistified air in distilled water impregnated with fixed air, than in plain distilled water, which is perfectly conformable to the abovementioned hypothesis; for as fixed air is a principal constituent part of vegetable substances, it is reasonable to suppose that the work of vegetation will go on better in water containing this substance, than in other water.

There are several memoirs of Mr. LAVOISIER published by the Academy of Sciences, in which he intirely discards phlogiston, and explains those phænomena which have been usually attributed to the loss or attraction of that substance, by the absorption or expulsion of dephlogistified air; and as not only the foregoing experiments, but most other phænomena of nature, seem explicable as well, or nearly as well, upon this as upon the commonly believed principle of phlogiston, it may be proper briefly to mention in what manner I would explain them on this principle, and why I have adhered to the other. In doing this, I shall not conform strictly to his theory, but shall make such additions and alterations as seem to suit it best to the phænomena; the more so, as the foregoing experiments may, perhaps, induce the author himself to think some such additions proper.

According to this hypothesis, we must suppose, that water consists of inflammable air united to dephlogistified air; that nitrous air, vitriolic acid air, and the phosphoric acid, are also combinations of phlogistified air, sulphur, and phosphorus, with dephlogistified air; and that the two former, by a further addition of the same substance, are reduced to the common
nitrous

nitrous and vitriolic acids; that the metallic calces consist of the metals themselves united to the same substance, commonly, however, with a mixture of fixed air; that on exposing the calces of the perfect metals to a sufficient heat, all the dephlogisticated air is driven off, and the calces are restored to their metallic form; but as the calces of the imperfect metals are vitrified by heat, instead of recovering the metallic form, it should seem as if all the dephlogisticated air could not be driven off from them by heat alone. In like manner, according to this hypothesis, the rationale of the production of dephlogisticated air from red precipitate is, that during the solution of the quicksilver in the acid and the subsequent calcination, the acid is decomposed, and quits part of its dephlogisticated air to the quicksilver, whereby it comes over in the form of nitrous air, and leaves the quicksilver behind united to dephlogisticated air, which, by a further increase of heat, is driven off, while the quicksilver re-assumes its metallic form. In procuring dephlogisticated air from nitre, the acid is also decomposed; but with this difference, that it suffers some of its dephlogisticated air to escape, while it remains united to the alkali itself, in the form of phlogisticated nitrous acid. As to the production of dephlogisticated air from plants, it may be said, that vegetable substances consist chiefly of various combinations of three different bases, one of which, when united to dephlogisticated air, forms water, another fixed air, and the third phlogisticated air; and that by means of vegetation each of these substances are decomposed, and yield their dephlogisticated air; and that in burning they again acquire dephlogisticated air, and are restored to their pristine form.

It seems, therefore, from what has been said, as if the phenomena of nature might be explained very well on this principle,

ciple, without the help of phlogiston ; and indeed, as adding dephlogisticated air to a body comes to the same thing as depriving it of its phlogiston and adding water to it, and as there are, perhaps, no bodies entirely destitute of water, and as I know no way by which phlogiston can be transferred from one body to another, without leaving it uncertain whether water is not at the same time transferred, it will be very difficult to determine by experiment which of these opinions is the truest ; but as the commonly received principle of phlogiston explains all phænomena, at least as well as Mr. LAVOISIER's, I have adhered to that. There is one circumstance also, which though it may appear to many not to have much force, I own has some weight with me ; it is, that as plants seem to draw their nourishment almost intirely from water and fixed and phlogisticated air, and are restored back to those substances by burning, it seems reasonable to conclude, that notwithstanding their infinite variety they consist almost intirely of various combinations of water and fixed and phlogisticated air, united according to one of these opinions to phlogiston, and deprived according to the other of dephlogisticated air ; so that, according to the latter opinion, the substance of a plant is less compounded than a mixture of those bodies into which it is resolved by burning ; and it is more reasonable to look for great variety in the more compound than in the more simple substance.

Another thing which Mr. LAVOISIER endeavours to prove is, that dephlogisticated air is the acidifying principle. From what has been explained it appears, that this is no more than saying, that acids lose their acidity by uniting to phlogiston, which with regard to the nitrous, vitriolic, phosphoric, and arsenical acids is certainly true. The same thing, I believe, may be said of the acid of sugar ; and Mr. LAVOISIER's experiment is a
strong

strong confirmation of BERGMAN's opinion, that none of the spirit of nitre enters into the composition of the acid, but that it only serves to deprive the sugar of part of its phlogiston. But as to the marine acid and acid of tartar, it does not appear that they are capable of losing their acidity by any union with phlogiston. It is to be remarked also, that the acids of sugar and tartar, and in all probability almost all the vegetable and animal acids, are by burning reduced to fixed and phlogisticated air, and water, and therefore contain more phlogiston, or less dephlogisticated air, than those three substances.

